

**Show Poultry Pre-Starter
MEDICATED**

FOR TURKEY AND BROILERS 0-4 WEEKS OF AGE

For improved feed efficiency and as aid in the prevention of coccidiosis.

ACTIVE DRUG INGREDIENT

Monensin	90 g/ton
Bacitracin Methylene Disalicylate	50 g/ton

GUARANTEED ANALYSIS

Crude Protein (Min)	30%
Lysine (Min)	1.7%
Methionine (Min)	.5%
Crude Fat (Min)	8%
Crude Fiber (Max)	4%
Calcium (Min)	1.1%
Calcium (Max)	1.5%
Phosphorus (Min)	.90%
Salt (Min)	.10%
Salt (Max)	.30%

INGREDIENTS

Grain Products, Plant Protein Products, Animal Protein Products, Forage Products, Processed Grain By-Products, Mono-calcium Phosphate, Di-calcium Phosphate, Calcium Carbonate, Roughage Products, Vitamin E Supplement, Vitamin A Supplement, Mineral Oil, Vitamin D3 Supplement, Silicon Dioxide, Choline Chloride, D-Calcium Pantothenate, Riboflavin Supplement, Niacin Supplement, Silicon Dioxide, Menadione Dimethylpyrimidinol Bisulfite, Biotin, Pyridoxine Hydrochloride, Thiamine Mononitrate, Folic Acid, Zinc Oxide, Condensed Lignin Sulfonate, Soybean Oil, Propionic Acid, Ferrous Sulfate, Manganous Oxide, Ascorbic Acid, Sodium Selenite, Copper Sulfate, Vitamin E Supplement, Vitamin B12 Supplement, Calcium Iodate, L-Lysine, Silicon Dioxide, Aluminum Oxide, Iron Oxide, Calcium Oxide, Potassium Oxide, Titanium Di-Oxide, DL-Methionine, Menadione Sodium Bisulfite Complex, Dried Fermentation Products, Verxite, Propionic Acid.

FEEDING DIRECTIONS

Feed continuously to growing broiler or turkeys as the sole ration from 1 day of age. The optimum level depends upon the severity of coccidiosis exposure.

Warning Statement: Do not feed to laying hens. Do not feed to chickens over 16 weeks of age. CAUTION:-----Do not allow horses, other equines, mature turkeys or guinea fowl access to feed containing Monensin. Ingestion of Monensin by horses, mature turkeys, and guinea fowl has been fatal. Some strains of turkey coccidian may be monensin tolerant or resistant

Manufactured By



22.6 Kg, 50 Pounds (Net)

F5039A