

# BABY BLOOMER 23% PIG PRE STARTER PELLETS MEDICATED

FOR PRE-STARTER PIGS WEIGHING 2 TO 11 POUNDS

Treatment of bacterial enteritis caused by Escherichia coli and Salmonella choleraesuis and bacterial pneumonia caused by Pasteurella multocida susceptible to Chlortetracycline.

For the control of swine dysentery associated with Brachyspira (formerly Serpulina or Treponema) hyodysenteriae susceptible to Tiamulin.

ACTIVE DRUG INGREDIENT	
Chlortetracycline	400 g/ton
Tiamulin Hydrogen Fumerate	35 g/ton

GUARANTEED ANALYSIS	
Crude Protein (Min)	23%
Lysine (Min)	1.2%
Crude Fat (Min)	6.25%
Crude Fiber (Max)	3.5%
Calcium (Min)	.70%
Calcium (Max)	1.1%
Phosphorus (Min)	.80%
Sodium (Min)	.25%
Sodium (Max)	.75%
Selenium (Min)	.3 ppm
Zinc (Min)	100 ppm

#### INGREDIENTS

Grain Products, Plant Protein Products, Animal Protein Products, Animal Plasma, Forage Products, Soybean Oil, Monocalcium Phosphate, Dicalcium Phosphate, Sugar, L-Lysine, D.L. Methionine, Calcium Carbonate, Dried Egg, Vitamin A Acetate with D-activated Animal Sterol (source of Vitamin D3), Vitamin E Supplement, Vitamin B12 Supplement, Riboflavin Supplement, Niacin Supplement, Calcium Pantothenate, Choline Chloride, Folic Acid, Menadione Sodium Bisulfite Complex, Pyridoxine Hydrochloride, Thiamine Mononitrate, d-Biotin, Dried Extracted Streptomyces Meal & Fermentation Solubles, Corn Distillers Dried Grains, Corn Distillers Dried Solubles & Condensed Fermented Corn Extractives, Ground Rice hulls, Sodium Selenite, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Ethylenediamine Dihydriodide, Artificial Flavors.

#### FEEDING DIRECTIONS

Start feeding to pigs when they are a few days old by placing a handful in the pen or in a shallow pan each day until the pigs are cleaning it up quickly. 50 pounds feed per litter is recommended.

WARNING: Withdraw 2 days before slaughter.

CAUTION: Do not feed CTC undiluted. Do not feed to animals other than swine. The effects of Tiamulin on swine reproductive performance, pregnancy and lactation have not been determined. Use as the only source of Tiamulin and Chlortetracycline. Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g. Monensin, Lasalocid, Narasin, Salinomycin and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feeds.

LIMITATIONS FOR USE: Feed continuously as sole ration for 14 days. Feed approximately 400 g/ton Chlortetracycline, varying with bodyweight and feed consumption to provide 10 mg/lb. of bodyweight per day. FDA approval provides only for use of Chlortetracycline products manufactured under NADA 046-699 (ChlorMax), 048-761 (Aureomycin) or 138-935 (Pennchlor).

Caution: Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

Manufactured By  
  
BEHREND'S  
FEED & FERTILIZER L.P.  
3599 Ranch Road 1376  
Fredericksburg, TX 78624  
830.997.3410

22.6 Kg, 50 Pounds (Net)

## F6200

## 3/16