

# 14% CREEP - (B) PELLETS MEDICATED

(For Ruminants Only)

THIS FEED IS DESIGNED TO BE FED TO STARTER LAMBS

For the prevention of coccidiosis caused by Eimeria ovina, E. Crandallis, E. ovinoidalis (E. ninakohlyakimovae), E. parva and E. intricata in sheep maintained in confinement.

## ACTIVE DRUG INGREDIENT

Lasalocid (as Lasalocid Sodium) 30 g/ton

## GUARANTEED ANALYSIS

Crude Protein (Min)	14%
Crude Fat (Min)	2%
Crude Fiber (Max)	12%
Calcium (Min)	.50%
Calcium (Max)	1.0%
Phosphorus (Min)	.35%
Salt (Min)	.40%
Salt (Max)	.60%
Selenium (Min)	.1 ppm
Vitamin A (Min)	3,000 IU/lb

## INGREDIENTS

Grain Products, Plant Protein Products, Roughage Products, Forage Products, Processed Grain By-Products, Ammonium Chloride, Calcium Carbonate, Monocalcium Phosphate, Dicalcium Phosphate, Salt, Roughage Products, Vitamin E Supplement, Vitamin A Supplement, Mineral Oil, Vitamin D3-Supplement, Silicon Dioxide, Zinc Oxide, Ferrous Carbonate, Manganous Oxide, Calcium Iodate, Sodium Selenite, Cobalt Carbonate, Molasses Products, Urea Formaldehyde Condensation Polymer and Silica.

## FEEDING DIRECTIONS

This is a complete creep feed for suckling lambs. Keep creep feed before lambs at all times once they are on full feed. Feed continuously 1 to 4.6 pounds of feed to provide not less than 15 mg nor 70 mg of Lasalocid per head per day depending on body weight. Large lambs which have not been on feed should be started on feed gradually. If it becomes necessary to withhold or limit feeding of animals on full feed, the return to full feeding must be gradual in order to avoid disturbances resulting from overeating.

**CAUTION:-----** The safety of Lasalocid for use in unapproved species has not been established. Do not allow horses or other equines access to feeds containing Lasalocid as ingestion may be fatal.

Manufactured By:



22.6 Kg, 50 Pounds (Net)

**F0124H**

**3/16**