

ZOO & WILDLIFE DEWORMER

FOR ZOO & WILDLIFE ANIMALS

For the removal and control of small stomach worms (*Trichostrongylus* spp.), thread-necked intestinal worms (*Nematodirus* spp.), barberpole worms (*Haemonchus* spp.) and whipworms (*Trichuris* spp.).

ACTIVE DRUG INGREDIENT

Fenbendazole 217.6 g/ton

GUARANTEED ANALYSIS

| | |
|---------------------|--------------|
| Crude Protein (Min) | 16% |
| Crude Fat (Min) | 2.5% |
| Crude Fiber (Max) | 15% |
| Calcium (Min) | .6% |
| Calcium (Max) | .9% |
| Phosphorus (Min) | .6% |
| Salt (Min) | .3% |
| Salt (Max) | .7% |
| Potassium (Min) | .9% |
| Vitamin A (Min) | 5,000 IU/ Lb |

INGREDIENTS

Grain Products, Plant Protein Products, Forage Products, Processed grain product, Roughage Products 8%, Molasses Products, Monocalcium Phosphate, Dicalcium Phosphate, Salt, Calcium Carbonate, Vitamin A Acetate with D-activated Animal Sterol (source of Vitamin D3), Vitamin E Supplement, Vitamin B12 Supplement, Riboflavin Supplement, Niacin Supplement, Calcium Pantothenate, Choline Chloride, Folic Acid, Menadione Sodium Bisulfite Complex, Pyridoxine Hydrochloride, Thiamine Mononitrate, d-Biotin, Manganous Oxide, Zinc Oxide, Ferrous Carbonate, Calcium Iodate, Cobalt Carbonate, Condensed Lignin Sulfonate, Propylene Glycol, Sodium Acetate, Sodium Sulfate, Artificial Flavors added.

FEEDING DIRECTIONS

Feed 2.5mg/kg of bodyweight per day for 3 days.

LIMITATION ----- Feed as per use level over a 3-day period. Retreatment may be needed after 6 weeks. Do not use 14 days before or during hunting season.

NOTE ----- Consult veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Manufactured For



**BEHREND'S FEED
& FERTILIZER LLC**
3599 Ranch Road 1376
Fredericksburg, TX 78624
(830)-997-3410

22.6 Kg, 50 Pounds (Net)

F1101Y

3/16