

Goat Feed 14% (D) Medicated

(For Ruminants Only)

THIS FEED IS DESIGNED TO BE FED TO GROWER GOATS

For the prevention of coccidiosis in young goats caused by Eimeria Christenseni and Eimeria Ninakohlyakimovae

ACTIVE DRUG INGREDIENT

Decoquinatate .003%

GUARANTEED ANALYSIS

Crude Protein (Min)	14%
(This includes not more than 1% equivalent protein from non-protein nitrogen.)	
Crude Fat (Min)	2.0%
Crude Fiber (Max)	22%
Calcium (Min)	.80%
Calcium (Max)	1.0%
Phosphorus (Min)	.25%
Salt (Min)	.30%
Salt (Max)	.70%
Selenium (Min)	.3 ppm
Vitamin A (Min)	3,000 IU/Lb

INGREDIENTS

Grain Products, Plant Protein Products, Forage Products, Roughage Products 18%, Molasses Products, Calcium Carbonate, Monocalcium Phosphate, Dicalcium Phosphate, Condensed Lignin Sulfonate, Salt, Ammonium Chloride, Vitamin A Supplement, Vitamin E Supplement, Mineral Oil, Vitamin D3 Supplement, Silicon Dioxide, Zinc Oxide, Ferrous Carbonate, Manganous Oxide, Calcium Iodate, Cobalt Carbonate, Sodium Selenite, Magnesium Oxide, Ethylenediamine Dihydriodide, Manganous Oxide, Verxite, Propionic Acid.

FEEDING DIRECTIONS

To be fed to young goats at a rate to provide 22.7 mg of Decoquinatate per 100 lbs of body weight daily- as indicated below.

FEEDING RATE

ANIMAL WEIGHT	PER HEAD PER DAY
30 lbs	0.5 lbs
60 lbs	1.0 lbs

Feed at least 28 days during period of exposure or when experience indicates that coccidiosis is likely to be a hazard

WARNING..... Do not feed to goats producing milk for food.

Manufactured By



**BEHREND'S FEED
& FERTILIZER LLC**
3599 Ranch Road 1376
Fredericksburg, TX 78624
(830)-997-3410

22.6 Kg, 50 Pounds (Net)

F3040A

3/16