

# Show Broiler Finisher Pellet MEDICATED

## COMPLETE FEED FOR BROILERS

For increased rate of weight gain and improved feed efficiency and aid in the prevention of coccidiosis.

### ACTIVE DRUG INGREDIENT

Bacitracin Methylene Disalicylate 50 g/ton  
Amprolium .0125 g/ton

### GUARANTEED ANALYSIS

Crude Protein (Min)	23%
Lysine (Min)	1.2%
Methionine (Min)	.50%
Crude Fat (Min)	7.0%
Crude Fiber (Max)	4.0%
Calcium (Min)	.60%
Calcium (Max)	.90%
Phosphorus (Min)	.45%
Salt (Min)	.10%
Salt (Max)	.30%

### INGREDIENTS

Grain Products, Plant Protein Products, Processed Grain By-Products, Soybean Oil, Animal Protein Products, Calcium Carbonate, Calcium Stearate, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Lecithin, Silicon Dioxide, Verite Granules, Citric Acid (a preservative), Butylated Hydroxytoluene (a preservative), Ethoxyquin (a preservative), Phosphoric Acid, Butylated Hydroxyanisole (a preservative), Bentonite, Diatomaceous Earth, Dry Kelp Meal, Yeast Culture, Active Dry Yeast, Natural Flavors, Basic Copper Chloride, Vitamin E Supplement, Mineral Oil, Manganese Hydroxychloride, Iron Sulfate, Niacin Supplement, Zinc Hydroxychloride, Manganese Sulfate, Manganous Oxide, Zinc Sulfate, Zinc Oxide, Dried Bacillus subtilis Fermentation Product, Sodium Sulfate, Hydrated Magnesium Silicate, Starch, Poly-vinyl Alcohol (PVA), Chromium Propionate, Biotin, Calcium Pantothenate, Wheat Flour, Potato Starch, Dried Trichoderma longibrachiatum Fermentation Extract, Vitamin B12 Supplement, Selenomethionine Hydroxy Analogue, Vanillin, Vitamin A Supplement, Vitamin D3 Supplement, Riboflavin Supplement, Menadione Nicotinamide Bisulfite, Pyridoxine Hydrochloride, 25-Hydroxyvitamin D3, Thiamine Mononitrate, Ethylenediamine Dihydroiodide, Folic Acid, Sodium Molybdate, Spray Dried Lignin Sulfate, Mono-calcium Phosphate, Di-calcium Phosphate, Roughage Products, L-Lysine, Salt, Choline Chloride, Dehydrated Cereal Grass and Corn Cobs, DL-Methionine, Sodium Bi-carbonate, Propionic Acid, Acetic Acid, Benzoic Acid, Sorbic Acid, Buffering Agents, L-Threonine, Wheat Flour, Dehydrated Pichia pastoris Fermentation Extract, Modified Starch.

### FEEDING DIRECTIONS

Feed as a sole ration to turkeys and/or broilers. Allow access to an adequate supply of fresh, clean water at all times.

WARNING: Use as sole source of Amprolium.

CAUTION: For replacement chicken only. Do not use AMPROLIUM in feeds containing bentonite.

Manufactured By



22.6 Kg, 50 Pounds (Net)

**F5036**

**3/16**