

Show Broiler Finisher MEDICATED

COMPLETE FEED FOR BROILERS

For improved feed efficiency and as aid in the prevention of coccidiosis.

ACTIVE DRUG INGREDIENT

Monensin	99 g/ton
Bacitracin Methylene Disalicylate	50 g/ton

GUARANTEED ANALYSIS

Crude Protein (Min)	21%
Lysine (Min)	1.1%
Methionine (Min)	.5%
Crude Fat (Min)	6%
Crude Fiber (Max)	4%
Calcium (Min)	1.0%
Calcium (Max)	1.4%
Phosphorus (Min)	.8%
Salt (Min)	.10%
Salt (Max)	.40%

INGREDIENTS

Grain Products, Plant Protein Products, Animal Protein Products, Forage Products, Mono-calcium Phosphate, Di-calcium Phosphate, Calcium Carbonate, Roughage Products, Vitamin E Supplement, Vitamin A Supplement, Mineral Oil, Vitamin D3 Supplement, Silicon Dioxide, Choline Chloride, D-Calcium Pantothenate, Riboflavin Supplement, Niacin Supplement, Silicon Dioxide, Menadione Dimethylpyrimidinol Bisulfite, Biotin, Pyridoxine Hydrochloride, Thiamine Mononitrate, Folic Acid, Zinc Oxide, Condensed Lignin Sulfonate, Soybean Oil, Ferrous Sulfate, Manganous Oxide, Ascorbic Acid, Sodium Selenite, Copper Sulfate, Vitamin B12 Supplement, Calcium Iodate, L-Lysine, Silicon Dioxide, Aluminum Oxide, Iron Oxide, Calcium Oxide, Potassium Oxide, Titanium Di-Oxide, DL-Methionine, Menadione Sodium Bisulfite Complex, Dried Fermentation Products, Verxite, Propionic Acid.

FEEDING DIRECTIONS

Feed continuously to finishing broilers as the sole ration. The optimum level depends upon the severity of coccidiosis exposure.

Warning Statement: Do not feed to laying hens. Do not feed to chickens over 16 weeks of age. CAUTION: -----For broiler chickens and replacement chickens intended for use as caged layers only. Do not allow horses, other equines, access to feed containing Monensin. Ingestion of Monensin by equines has been fatal

Manufactured By



**BEHREND'S FEED
& FERTILIZER LLC**
3599 Ranch Road 1376
Fredericksburg, TX 78624
(830)-997-3410

22.6 Kg, 50 Pounds (Net)

F5055C

3/16 and Crumbles