

BROILER FINISHER 21 CRUMBLES (R) MEDICATED

FINISHER BROILERS 5 WEEKS OF AGE TO MARKET

As an aid in the prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

ACTIVE DRUG INGREDIENT

Monensin 99 g/ton

GUARANTEED ANALYSIS

Crude Protein (Min)	21%
Lysine (Min)	1.1%
Methionine (Min)	.50%
Crude Fat (Min)	6%
Crude Fiber (Max)	4%
Calcium (Min)	1.0%
Calcium (Max)	1.4%
Phosphorus (Min)	.8%
Sodium (Min)	.10%
Sodium (Max)	.40%

INGREDIENTS

Grain Products, Plant Protein Products, Animal Protein Products, Forage Products, Processed Grain By-Products, Hydrolyzed Animal & Vegetable Fat, Monocalcium Phosphate, Dicalcium Phosphate, Calcium Carbonate, Roughage Products, Vitamin A Supplement, Vitamin E Supplement, Vitamin B12 Supplement, Riboflavin Supplement, Niacin Supplement, Mineral Oil, D-Calcium Pantothenate, Choline Chloride, Silicon Dioxide, Menadione Sodium Bisulfite Complex, Folic Acid, Pyridoxine Hydrochloride, Thiamine Mononitrate, d-Biotin, Vitamin D3 Supplement, Choline Chloride, Ferrous Sulfate, Manganous Oxide, Ascorbic Acid, Sodium Selenite, Calcium Iodate, Zinc Oxide, Copper Sulfate, L-Lysine, DL-Methionine, Urea Formadlehyde Condensation Polymer, Silica and Natural and Artificial Flavors.

FEEDING DIRECTIONS

Feed continuously as the sole ration.

WARNING: ----- Do not feed to laying chickens.

CAUTION: ----- Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by equines has been fatal. In the absence of coccidiosis, the use of Monensin with no withdrawal period may limit feed intake resulting in reduced weight gains.

Manufactured By



22.6 Kg, 50 Pounds (Net)

F5055C