

Show Turkey Grower Pellet MEDICATED

COMPLETE FEED FOR GROWING TURKEYS

For prevention of coccidiosis caused by Eimeria adenoeides, Eimeria meleagrimitis and Eimeria gallopavoni and for increased rate of weight gain, and improved feed efficiency.

ACTIVE DRUG INGREDIENT

Monensin	72 g/ton
Bacitracin Methylene Disalicylate	50 g/ton

GUARANTEED ANALYSIS

Crude Protein (Min)	23%
Lysine (Min)	1.2%
Methionine (Min)	.4%
Crude Fat (Min)	5.5%
Crude Fiber (Max)	4%
Calcium (Min)	1.0%
Calcium (Max)	1.4%
Phosphorus (Min)	.8%
Salt (Min)	.10%
Salt (Max)	.40%

INGREDIENTS

Grain Products, Plant Protein Products, Animal Protein Products, Forage Products, Mono-calcium Phosphate, Salt, Calcium Carbonate, Roughage Products, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Choline Chloride, D-Calcium Pantothenate, Riboflavin Supplement, Niacin Supplement, Silicon Dioxide, Biotin, Pyridoxine Hydrochloride, Thiamine Mononitrate, Folic Acid, Zinc Oxide, Condensed Lignin Sulfonate, Soybean Oil, Ferrous Sulfate, Manganous Oxide, Ascorbic Acid, Sodium Selenite, Copper Sulfate, Vitamin B12 Supplement, Calcium Iodate, L-Lysine, Silicon Dioxide, Aluminum Oxide, Iron Oxide, Calcium Oxide, Potassium Oxide, Titanium Di-Oxide, DL-Methionine, Menadione Sodium Bisulfite Complex, Dried Fermentation Products, Verxite, Propionic Acid.

FEEDING DIRECTIONS

Feed continuously to growing turkeys as the sole ration. The optimum level depends upon the severity of coccidiosis exposure.

CAUTION: -----Do not allow horses, other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Some strains of turkey coccidia may be Monensin tolerant or resistant. Monensin may interfere with development of immunity to turkey coccidiosis.

Manufactured By



22.6 Kg, 50 Pounds (Net)

F5060

3/16